



\*\*This news release from K-State Research and Extension is available online at <https://ksre-learn.com/warm-season-grass-calendar>

Released: March 21, 2025

### **Lawn Calendar: Warm-season grasses in Kansas**

K-State horticulture expert outlines tips for maintaining warm-season lawns

*K-State Research and Extension news service*

MANHATTAN, Kan. — While cool-season grasses are more common in Kansas, many parts of the state are more ideal for warm-season grasses, or those more tolerant of high temperatures and drought conditions.

Zoysiagrass, bermudagrass and buffalograss are the most common warm-season grasses in Kansas. Of these, buffalograss is thought to require less maintenance because it can survive with less water and fertilizer than the other two varieties.

The following calendar – provided by K-State horticulture expert Cynthia Domenghini -- will give homeowners a good plan for maintaining warm-season grasses in Kansas.

#### **March**

Zoysiagrass and bermudagrass – Spot treat broadleaf weeds, if necessary. Treat on a day that is 50 degrees Fahrenheit or warmer. Rain or irrigation within 24 hours of application will reduce its effectiveness.

Buffalograss – Spot treat broadleaf weeds, if necessary; fall treatments are more effective. Spray early enough in March that the buffalograss is still dormant. Treat on a day that is 50 degrees F or warmer. Rain or irrigation within 24 hours of application will reduce its effectiveness.

#### **April**

Zoysiagrass and bermudagrass – Apply crabgrass preventer between April 1 and 15, or apply preventer when the eastern redbud is approaching full bloom. Crabgrass preventers must be watered in before they will start to work.

Buffalograss – Apply crabgrass preventer between April 1 and 15 or when the eastern redbud is in full bloom. Crabgrass preventers must be watered in before they will start to work. Avoid using broadleaf herbicides as the buffalograss is greening up, which may cause injury.

### **May through August 15**

Zoysiagrass and bermudagrass – Fertilize with one pound of nitrogen per 1,000 square feet. Follow recommendations on the bag. More applications will give a deeper green color, but will increase mowing and lead to thatch buildup in zoysiagrass. Bermudagrass can have problems with thatch buildup, but is less likely.

For bermudagrass, consider 2 to 4 applications. For zoysiagrass, consider 1 to 2 applications.

Depending on what you decide, the suggested timeframe for applications includes:

One application – apply in June.

Two applications – apply in May and July.

Three applications – apply in May, June and early August.

Four applications – apply in May, June, July and early August.

June

Zoysiagrass and bermudagrass – If grubs have been a problem in the past, apply a product containing imidacloprid (May through June) or chlorantraniliprole (May). These products must be watered in before they are effective. June is a good time to core aerate a warm-season lawn.

Buffalograss – Fertilize with 1 pound of nitrogen per 1,000 square feet. If desired, fertilize again in July. If grubs have been a problem in the past, apply a product containing imidacloprid (May through June) or chlorantraniliprole (May). These products can also be applied in mid-May if there are problems with billbugs or May beetle grubs.

### **Late July through August**

Zoysiagrass and bermudagrass – Apply a grub killer if you see grub damage. If imidacloprid has been applied previously, this should not be necessary. Grub killers must be watered in immediately.

Buffalograss – Apply a grub killer if you see grub damage. If imidacloprid has been applied previously, or if grubs have not been a problem in the past, this should not be necessary. Grub killers must be watered in immediately.

### **Late October**

Zoysiagrass and bermudagrass – Spray for broadleaf weeds if they are a problem. Treat on a day that is at least 50 degrees F. Rain or irrigation within 24 hours reduces the effectiveness of the spray. Use the rates listed on the product label.

Buffalograss – Spray for broadleaf weeds if they are a problem. Look carefully because such winter annuals as chickweed and henbit are small and easily overlooked. Work with your local extension agencies for control recommendations. Treat on a day that is at least 50 degrees F. Rain or irrigation within 24 hours reduces the effectiveness of the spray. Use the rates listed on the product label.

“Now is a great time to get the soil tested to determine nutrient levels,” Domenghini said. “The best start to a healthy lawn is healthy soil. Wait to apply fertilizers until you know what your turfgrass needs. This step can save homeowners time and money.”

Domenghini and her colleagues in K-State's Department of Horticulture and Natural Resources produce a [weekly Horticulture Newsletter](#) with tips for maintaining home landscapes and gardens.

Interested persons can subscribe to the newsletter, as well as send their garden and yard-related questions to [hortsupport@ksu.edu](mailto:hortsupport@ksu.edu), or contact your [local K-State Research and Extension office](#).

-30-

**FOR PRINT PUBLICATIONS:** Links used in this story  
K-State Horticulture Newsletter, <https://hnr.k-state.edu/extension/info-center/newsletters/index.html>

K-State Research and Extension local offices, <https://www.ksre.k-state.edu/about/statewide-locations.html>

K-State Research and Extension is a short name for the Kansas State University Agricultural Experiment Station and Cooperative Extension Service, a program designed to generate and distribute useful knowledge for the wellbeing of Kansans. Supported by county, state, federal and private funds, the program has county extension offices, experiment fields, area extension offices and regional research centers statewide. Its headquarters is on the K-State campus in Manhattan. For more information, visit [www.ksre.ksu.edu](http://www.ksre.ksu.edu). K-State Research and Extension is an equal opportunity provider and employer.

**For more information:**  
Cynthia Domenghini  
[Cdom@ksu.edu](mailto:Cdom@ksu.edu)